

Issue 1, Volume 2, February 2009  
Scandinavian Journal of Optometry and Visual Science

# SJOVS

## Refractive trends in Swedish Military recruits



■ Her presenteres forskningsarbeider som tilfredsstillter kriteriene for trykking av vitenskapelige artikler, utarbeidet av International Committee of Medical Journals Editors (ICMJE). "Uniform requirements for manuscripts submitted to biomedical journals: writing and editing for biomedical publication" kan leses på <http://www.icmje.org/>. Utdrag av kriteriene kan leses på norsk på [www.optikerne.no/optikeren](http://www.optikerne.no/optikeren). En objektiv, uavhengig og kritisk vurdering foretatt av en fagfelle, er vesentlig for de vitenskapelige arbeidene som presenteres.

■ Research papers presented in this part of our journal fulfill the criteria for the printing of scientific articles made by the International Committee of Medical Journals Editors (ICMJE). "Uniform requirements for manuscripts submitted to biomedical journals: writing and editing for biomedical publication" can be read at <http://www.icmje.org/>. Unbiased, independent, critical assessment through peer review is an important part of the scientific papers which are presented.

**Redaksjon**

Inger Lewandowski  
(Ansvarlig redaktør/Managing Editor)

**Anneli Demberg**

(Vitenskapelig redaktør/Editor-in-Chief)

Anneli Demberg, BSc, er PhD-student ved Glasgow Caledonian University, Skottland. Hun har tidligere vært ansatt som ved Avdeling for optometri og synsvitenskap. Hennes forskning heter: Childhood Amblyopia, development of a test protocol for identifying functional losses in childhood amblyopia & monitoring treatment efficacy.

Mailadresse: [sjovs@optikerforbund.no](mailto:sjovs@optikerforbund.no)

**Assosierte redaktører**

(Associate Editors)

**Rigmor C. Baraas**, BSc (Hons), PhD, 1. amanuensis, Avdeling for optometri og synsvitenskap, Høgskolen i Buskerud

**Per O. Lundmark**, BSc (Hons), PhD, 1. amanuensis, Avdeling for optometri og synsvitenskap, Høgskolen i Buskerud

**Vibeke Sundling**, BSc (Hons), MSc, høgskolelektor/stipendiat, Avdeling for optometri og synsvitenskap, Høgskolen i Buskerud

**Redaksjonskomité**

(Board of Editors)

**Leiv S. Bakketeig**, øyelege dr. med., overlege, professor ved Statens Institutt for Folkehelse, Oslo

**Rune L. Brautaset**, Mphil, PhD, BSc, MPhil, PhD, Programdirektør optometriutdanningen, St Eriks Ögonsjukhus, Karolinska Institutet, Stockholm, Sverige

**Jan Richard Bruenech** BSc (Hons), PhD, professor, Avdeling for optometri og synsvitenskap, Høgskolen i Buskerud, Kongsberg og professor II ved Randers Tekniske Høyskole, Danmark

**Liv Drolsum**, øyelege dr. med., professor II, Øyeavdelingen, Ullevål Universitetssykehus, Oslo

**Jörgen Gustafsson**, PhD, forskningsleder ved Avdeling for optometri, Högskolan i Kalmar, Sverige

**Gunnar Horgen** PhD, professor, prodekan, Avdeling for optometri og synsvitenskap, Høgskolen i Buskerud, Kongsberg

**Internasjonale rådgivere**

(International advisory board):

**Feike Grit**, Holland

**Svend-Erik Runberg**, Danmark

Det er med stor glede og stolthet at vi i dette nummeret av Optikerens presenterer vår første fagfelleverderte artikkel. Det er en kort, men interessant forskningsrapport om synet hos svenske rekrutter.

Da vi introduserte SJOVS for våre lesere for over ett år siden, hadde vi nok trodd at etableringen av en slik fagdel skulle være enklere enn den har vært. Vi har imidlertid godt håp om at det nå etter hvert vil komme inn flere artikler, og selv om godkjenningprosessen tar tid, håper vi å kunne presentere flere artikler dette året.

Samtidig vil vi oppfordre de av våre lesere som har gjennomført kasus eller undersøkelser om å vurdere innsending av manuskriptet til SJOVS. For å kunne bli antatt må artikler tilfredsstillende visse krav til hvordan de skal skrives. Slå opp på Optikerens hjemmeside, som du finner ved å gå inn på NOFs hjemmeside: [www.optikerne.no](http://www.optikerne.no) Klikk på bildet av Optikerens på høyre side i skjermbildet. Under "vitenskapelige artikler" finnes informasjon om hva som kreves av artikkelforfattere og hva fagfellevurderingen går ut på.

*Vi minner om SJOVS' målsetting:*

SJOVS vurderer artikler som omhandler enten grunnleggende aspekter innen synsvitenskap, eller kliniske og anvendte tema som er relevante for utførelse og

anvendelse av optometri. Materialet kan være av eksperimentell, observerende eller historisk art. Emner som er viktige for optometristere vektlegges, men mange av temaene er også av interesse for psykologer, fysiologer og andre som er opptatt av utvikling, bruk og rehabilitering av syn. Grunnleggende studier kan omhandle utvikling og bruk av synet. Artikler som tar for seg utførelsen av optometri og instrumenter relatert til dette, utgjør en stor del av tidsskriftet, tema spenner fra design av brilleinnfatninger og kontaktlinser til øyesykdommer og bruk av medikamenter. Problemer med bruk av synet i bestemte situasjoner vektlegges også, for eksempel syn ved bilkjøring, bruk av optiske instrumenter og skjermarbeid.

Anneli Demberg

Vitenskapelig redaktør

Inger Lewandowski

Ansvarlig redaktør

# Refractive trends in Swedish Military recruits

D. Uhlin<sup>1</sup>, S. Lutteman<sup>1</sup>, J.A.M. Jennings<sup>2</sup>, R.L. Brautaset<sup>1</sup>

<sup>1</sup>Unit of Optometry, Department of Clinical Science, Karolinska Institutet, Box 8056, 104 20 Stockholm, Sweden and <sup>2</sup>Faculty of Life Science, University of Manchester, PO Box 88, Manchester M60 1QD, UK

## Abstract

**Purpose:** The aim of the study was to determine the prevalence of refractive errors in military recruits, with special reference to myopia.

**Methods:** 651 male military recruits, 17-23 years of age, were refracted at Pliktverket in Näsby (a National Service Administration office near Stockholm, Sweden). The refractive measurements were taken with a Topcon KR-7000P auto keratometer during the initial health examination of National Service recruits.

**Results:** Out of the 651 recruits the prevalence of myopia ( $\geq -0.50$ DS) was 37.7% (i.e., both eyes being myopic). The mean refractive error was RE  $-0.47$ DS and LE  $-0.41$ DS. This difference was not statistically significant.

**Discussion:** The prevalence of myopia in Swedish military recruits has increased from 8.9% to 37.7% and the mean refractive error has changed from  $+0.53$ DS to  $-0.44$ DS since the study of Strömberg (1936). However, the prevalence of myopia in these present day recruits is lower than the 49.5% found in 12-13 year old Swedes (Villarreal et al., 2000).

**Key words:** myopia – prevalence  
– Swedish male military recruits

## Correspondence:

Rune L. Brautaset  
Unit of Optometry  
Department of Clinical Science  
Karolinska Institutet  
Box 8056  
S-104 20 Stockholm  
Sweden  
Fax: +46 (0)8 672 3846  
E-mail: rlb@ste.ki.se

**Received:** February 2008

**Revised form:** November 2008

**Accepted:** December 2008

## Sammanfattning

**Bakgrund:** Målsättningen med studien var att undersöka prevalensen av synfel hos mönstrande män med fokus på myopiprevalens.

**Metod:** 651 värnpliktiga i åldern 17 till 23 år blev undersökta vid mönstring hos Pliktverket i Näsby utanför Stockholm. Mätningarna utfördes med en Topcon autorefraktor (KR-7000P) som del av deras hälsokontroll.

**Resultat:** Av de 651 värnpliktiga som undersöktes påträffades en myopiprevalens ( $\geq -0.50$ DS) på 37,7%. Det genomsnittliga synfelet var HÖ  $-0,47$ DS och VÖ  $-0,41$ D, en skillnad som inte var statistisk signifikant.

**Diskussion:** Myopiprevalensen hos mönstrande män i Sverige har ökat från 8,9% till 37,7% och det genomsnittliga synfelet har ändrat sig från  $+0,53$ D till  $-0,44$ DS sedan motsvarande studie genomfördes av Strömberg (1936). Samtidigt ser man att myopiprevalensen hos dagens mönstrande män är lägre än myopiprevalensen (49,5%) som hittades hos svenska 12-13 år gamla barn (Villarreal et al., 2000).

## Introduction

There is general agreement that the environment and genetics interact in the development of myopia (Adams and McBrien, 1992; Chung et al., 1996; Cordain et al., 2002; Mutti et al., 1996; O'Hara and Nelson, 1997) and that some populations, particularly in the Far East, are markedly predisposed to myopia (Hosaka, 1988; Lin et al., 1988; Saw et al., 2005; Wong et al., 2000; Zhao et al., 2000), but there is also evidence of increased prevalence of myopia in other parts of the world (Bar Dayan et al., 2005). The theory of myopia as an environmentally induced physiological adaptation to near work has lately been gaining more support (Adams and McBrien, 1992; Cordain et al., 2002; Flitcroft, 1998; Kinge and Midelfart, 1994; Kinge et al., 1998, 2000; Nyman 1998; Midelfart, 1990; Midelfart et al., 1992; Mutti et al., 2002; Saw et al., 2002; Williams et al., 2008), while the Sydney Myopia study recently reported that near work was not associated with concurrent myopia (Ip et al., 2007). From twin studies evidence has been presented that indicates that myopia is 80-90% heritable (Hammond et al., 2001; Lyhne, et al., 2001), however, family aggregation studies suggest lower heritability (Framingham Offspring Eye Study Group, 1996). Furthermore, the observed large changes in prevalence over very few generations have again illustrated the importance of environmental

factors (Morgan and Rose, 2002). In general, recent research can be summarized by saying that both environmental and genetic factors are of importance in the development of myopia, but that it is still to be discovered how these factors interact in order to explain the increase in myopia that can be seen in different parts of the world.

In the Nordic countries the prevalence of myopia is reported to be approximately 30% (Fledelius, 1983; Fledelius, 2000; Kinge et al., 1998; Midelfart, 2004). A summary of Nordic studies into the prevalence of myopia can be found in Table 1.

In 1936 Strömberg, based on subjective refraction, found the prevalence of myopia in 2616 Swedish conscripts aged about 20 years to be 8.9%. Since the study of Strömberg (1936) no study has been conducted in Sweden in order to determine the prevalence of myopia in a population of similar age. The aim of the current study was to determine the prevalence of today's Swedish army recruits and to compare the results with those of Strömberg (1936).

**Table 1:** Prevalence of myopia as found in the Nordic countries

Country	Author	Definition	Age group	Prevalence of myopia
Norway	Holst & Tjåland (1962)	?	9 years	2.2%
	Midelfart et al. (1992)	$\geq -0.25$	12 years	5.9%
	Midelfart et al. (2004)	$\geq -0.50$	Medical students	50.3%
			20-25 years	35%
Denmark	Øster & Kjærgaard* (1964)	$\geq -0.25$	7-8 years	1%
	Engbæk* (1970)	$\geq -0.25$	7-8 years	1.2%
	Johansen* (1950)	$\geq -0.25$	9-11 years	7.4%
	Fledelius (1976)	$\geq -0.25$	9-11 years	9.3%
	Jensen (1991)	$\geq -0.25$	7-12 years	28.6%
	Øster & Kjærgaard* (1964)	$\geq -0.25$	13-14 years	10%
	Goldschmidt (1968)	$\geq -0.25$	13-14 years	9.5%
	Engbæk* (1983)	$\geq -0.25$	13-14 years	11-12%
	Goldschmidt (1968)	$\geq -0.25$	17-23 years	14.5%
	Fledelius (1980)	$\geq -0.25$	17-23 years	13-14%
	Fledelius (1983)	$\geq -0.25$	17-23 years	32.6%
	Fledelius (200)	$\geq -0.50$	22-41 years (median 26 years)	50.0%
	Finland	Heinonen (1934)	$\geq -0.5$	14-15 years
Laatikainen & Erkkilä (1980)		$\geq -0.5$	7-8 years	1.9%
			11-12 years	7.2%
			14-15 years	21.8%
Mäntyjärvi (1983)		$\geq -0.25$	12 years	14%
			15 years	22.8%
Sweden	Villarreal et al. (2000)	$\geq -0.5$	12-13 years	49.7%
	Lundgren (1954)	?	18-19 years	19.5 - 25.3%
	Strömberg (1936)	$\geq -0.25$ $\geq -0.75$ $> -1.0$	20 years	8.9%
			(all men)	6.4%
				4.7%
				27%
	Stenström (1947)	$\geq -0.25$ $> -1.0$	20-35 years	16%
	Present study	$\geq -0.5$	17-23 years (all men)	37.7%

## Subjects and Methods

A total of 651 men aged between 17 and 23 years (mean age 18 years) was examined. The ocular examination was an additional and voluntary part of the initial health examination for National Service recruits. The examination took place at Pliktverket in Näsby (a National Service Administration office near Stockholm, Sweden). All recruits enrolled over a one week period were examined consecutively, all recruits accepted to be part of the study and none of the enrolled recruits were excluded. The recruits came from the greater Stockholm area. Of the 651 tested 98.5% were of Caucasian origin. All subjects gave informed consent.

All subjects had their visual acuity measured with a 4 metre LogMar chart. The refractive error was determined by use of a Topcon KR-700OP (auto kerato-refractometer) without the use of cycloplegics. As a safeguard against proximal effects prior to the auto-refraction all subjects were asked to fixate an acuity chart placed 4 metres away in order to relax accommodation.

All statistics were performed using InStatTM (GraphPad, La Jolla, CA, USA) and OriginTM (Origin Lab, Northampton, MA, USA) statistical software. Comparison of the present data and the results of Strömberg (1936) was done using Chi-squared test. The range of refractive errors could be considered normally distributed and the 95% confidence interval of the mean was constructed using the Student T-test.

## Results

Classification of refractive error was based on spherical equivalent (spherical value combined with half the cylinder value) and was divided into three groups: myopia ( $\geq -0.50D$ ), hyperopia ( $\geq +0.50D$ ), and emmetropia ( $+0.50D < \text{spherical equivalent} < -0.50D$ ).

*Myopia:* of the 651 recruits examined the overall prevalence of ( $\geq -0.50D$ ) was 37.7%. The prevalence of right eye myopia was 39%.

*Hyperopia:* ( $>+0.50D$ ) the overall prevalence was 19.6% and right eye prevalence was 17.8%.

*Emmetropia:* an overall prevalence of 42.7% with a 43.2% prevalence in the right eye.


The mean refractive error for the right eye was  $-0.47D$  ( $\pm 0.14$ ; 0.95 confidence interval of the mean) and for the left eye  $-0.41D$  ( $\pm 0.14$ ; 0.95 confidence interval of the mean).

The highest refractive errors were  $-8.25D$  and  $+6.50D$ . The distribution of refractive errors can be seen in Figure 1.

## Discussion

The subject group examined in this study is similar to the group examined by Strömberg (1936) in that they are all male, of similar age, and predominantly Caucasian (98.5%). This implies that the prevalence of myopia over a three-generation period (65 years) has increased from 8.9% to 37.7% ( $p < 0.05$ ), and the mean refractive error has changed from  $+0.53DS$

Figure 1: Distribution of refractive error (spherical equivalent) in the right and left eye



to -0.44DS ( $p < 0.05$ ). However, there are several factors that should be kept in mind when comparing the present study with the data of Strömberg (1936). Strömberg's data are based on subjective refraction without the use of cycloplegics whereas the present data are based on auto-refraction without the use of cycloplegic drugs. Both methods may overestimate the prevalence of myopia since both lack proper control of accommodation. However, Midelfart et al. (1992) found no significant difference between the spherical equivalent measured with an auto-refractor and that measured with subjective refraction, something that implies that the results are comparable.

The aim of the present study was to determine the prevalence of myopia in today's Swedish army recruits and to compare the results with the results of Strömberg (1936), and not to determine any cause for the development and/or increase in myopia. However, large changes in "normal life" have happened since Strömberg's time. In 1936 normal education lasted for 6 to 9 years. For the subjects in this study the average number of years of education was about 12 years. In addition to the increased reading implicit in extended education the popular present day near vision activities such as watching TV, video games and computers did not exist at the time of Strömberg.

It is interesting to note that over the same time period the average height of Swedish 18 year old men enrolled for National Service has increased by 2.3% (from 175.9 to 179.9 cm), while the average weight has increased by 22.1% (from 60.3 to 73.6 kg). This is an increase in body mass index (BMI) of 16.5% and is presumably a reflection of an increasingly sedentary lifestyle. (Height and weight data is extrapolated from statistical material published by Pliktverket (2002).)

The prevalence of myopia (49.5%) found by Villarreal et al. (2000) among Swedish 12-13 years old teenagers is much higher than our results for 17-23 year olds and those from other Nordic countries. It remains to be seen if this presages increasingly high levels of myopia in future generations of Swedes.

### Acknowledgement

The authors would like to thank the staff at Pliktverket for all their help in making this investigation possible. All the raw data was collected and summarized by Mr. Uhlin as part of his BSc dissertation.

### References

- Adams, D. W. and McBrien, N. A. (1992). Prevalence of myopia and myopic progression in a population of clinical microscopists. *Optom Vis Sci*, **69**, 467-473
- Bar Dayan, Y., Levin, A., Morad, Y., Grotto, I., Ben-David, R., Goldberg, A., Onn, E., Avni, I., Levy, Y. and Benyamini, O. G. (2005). The changing prevalence of myopia in young adults: a 13-year series of population-based prevalence surveys. *Invest Ophthalmol Vis Sci*, **46**, 760-2765
- Chung, K. M., Mohidin, N., Yeow, P. T., Tan, L. L. and O'Leary, D. (1996). Prevalence of visual disorders in Chinese schoolchildren. *Optom Vis Sci*, **73**, 695-700
- Cordain, L., Eaton, S. B., Miller, J. B., Lindeberg, S. and Jensen, C. (2002). An evolutionary analysis of the aetiology and pathogenesis of juvenile-onset myopia. *Acta Ophthalmol Scand*, **80**, 125-135
- Fledelius, H. (1976). Prematurity and the eye. *Acta Ophthalmol (Suppl)*, **128**, 3-245
- Fledelius, H. C. (1983). Is myopia getting more frequent? A cross sectional-study of 1416 Danes aged 16 years +. *Acta Ophthalmol (Copenh)*, **61**, 545-559
- Fledelius, H. C. (1988). Myopia prevalence in Scandinavia. A survey, with emphasis on factors of relevance for epidemiological refraction studies in general. *Acta Ophthalmol (Suppl)*, **185**, 44-50
- Fledelius, H. C. (2000). Myopia profile in Copenhagen medical students 1996-98. Refractive stability over a century is suggested. *Acta Ophthalmol Scand*, **78**, 501-505
- Flitcroft, D. I. (1998). Ophthalmologists should consider the cause of myopia and not simply treat its consequences. *Br J Ophthalmol*, **82**, 210-211
- Framingham Offspring Eye Study Group (1996). Familial aggregation and prevalence of myopia in the Framingham Offspring Eye Study. *Arch Ophthalmol*, **114**, 326-332
- Goldschmidt E (1968): On the etiology of myopia. An epidemiological study. *Acta Ophthalmol (Suppl)*, **98**, 11-172
- Hammond, C. J., Snieder, H., Gilbert, C. E. and Spector, T. D. (2001). Genes and environment in refractive error: the twin eye study. *Invest Ophthalmol Vis Sci*, **42**, 1232-1236
- Heinonen, E. (1934). Weiter Studien über die Schulmyopie. *Acta Ophthalmol Scand*, **12**, 110-121
- Holst, J. C. and Tjåland, J. (1962). Noen tall fra skoleøyelegekontoret i Oslo. *Tidsskr Nor Legeforen*, **19**, 1291-1294
- Hosaka, A. (1988). Population studies – Myopia experience in Japan. *Acta Ophthalmol (Suppl)*, **185**, 37-40
- Ip, J. M., Huynh, S. C., Robaei, D., Rose, K.A., Morgan, J. G., Smith, W., Kiffey, A. and Mitchell, P. (2007). Ethnic differences in the impact of parental myopia: findings from a population-based study of 12-year-old Australian children. *Invest Ophthalmol Vis Sci*, **48**, 2520-2528
- Jensen H (1991): Myopia progression in young school children. A prospective study of myopia progression and the efficacy of a trial with bifocal lenses and beta blocker eye drops. *Acta Ophthalmol (Suppl)*, **200**, 1-79

- Kinge, B. and Midelfart, A. (1994). Refractive errors among engineering students in Norway. *Ophthalmic Epidemiol*, **1**, 5-13
- Kinge, B., Midelfart, A. and Jacobsen, G. (1998). Refractive errors among young adults and university students in Norway. *Acta Ophthalmol Scand*, **76**, 692-695
- Kinge, B., Midelfart, A., Jacobsen, G. and Rystad, J. (2000). The influence of near-work on development of myopia among university students. A three-year longitudinal study among engineer students in Norway. *Acta Ophthalmol Scand*, **78**, 26-29
- Laatikainen, L. and Erkkilä, H. (1980). Refractive errors and the findings in school children. *Acta Ophthalmol (Copenh)*, **58**, 129-136
- Lin, L. K., Chin, C. J., Hung, P. T. and Ko, L. S. (1988). Nation-wide survey of myopia among schoolchildren in Taiwan. *Acta Ophthalmol (Suppl)*, **185**, 29-33
- Lundgren, P. G. (1954). Myopie in den höheren Lehranstalten Schwedens. *Klin Monatsbl Augenhkd*, **124**, 110
- Lyhne, N., Sjolie, A. K., Kyvik, K. O. and Green, A. (2001). The importance of genes and environment for ocular refraction and its determiners: a population based study among 20-45 year old twins. *Br J Ophthalmol*, **85**, 1470-1476
- Midelfart, A. (1990). Myopia and radial keratotomy: a survey among Norwegian ophthalmologists. *Acta Ophthalmol (Copenh)*, **68**, 597-600
- Midelfart, A., Aamo, B., Sjøhaug, K. A. and Dysthe, B. E. (1992). Myopia among medical students in Norway. *Acta Ophthalmol (Copenh)*, **70**, 317-322
- Midelfart A, Kinge B, Midelfart S & Lydersen S. (2004). Prevalens av brytningsfeil i Norge. *Tidsskr Nor Lægeforen*, **124**, 46-48
- Morgan, I. and Rose, K. (2005). How genetic is school myopia? *Prog Retin Eye Res*, **24**, 1-38
- Mutti, D. O., Zadnik, K. and Adams, A. J. (1996). Myopia. The nature versus nurture debate goes on. *Invest Ophthalmol Vis Sci*, **37**, 952-957
- Mutti, D. O., Mitchell, G. L., Moeschberger, M. L., Jones, L. A. and Zadnik, K. (2002). Parental myopia, near work, school achievement, and children's refractive error. *Invest Ophthalmol Vis Sci*, **43**, 3633-3640
- Mäntyjärvi, M. (1983). Incidence of myopia in a population of Finnish school children. *Acta Ophthalmol (Copenh)*, **61**, 417-423
- Nyman, K. G. (1988). Occupational near-work myopia. *Acta Ophthalmol (Suppl)*, **185**, 167-171
- O'Hara, M. A. and Nelson, L. B. (1997). Inheritance of myopia. In: Lippincott (ed.) *Duane's Ophthalmology*, Lippincott-Raven, London, Vol 3
- Plikverket (2002). The National Service Administration's internet publications (<http://www.pliktverket.se>)
- Saw, S. M., Chua, W. H., Hong, C. Y., Wu, H. M., Chan, W. Y., Chia, K. S., Stone, R. A. and Tan, D. (2002). *Nearwork in early-onset myopia. Invest Ophthalmol Vis Sci*, **43**, 332-339
- Saw, S. M., Tong, L., Chua, W. H., Chia, K. S., Koh, D., Tan, D. T. and Katz, J. (2005). Incidence and progression of myopia in Singaporean school children. *Invest Ophthalmol Vis Sci*, **46**, 51-57
- Stenström, S. (1947). Untersuchungen über die Variation und Kovariation der optischen Elemente des menschlichen Auges. *Acta Ophthalmol (Suppl)*, **26**
- Villarreal, M. G., Ohlsson, J., Abrahamsson, M., Sjöström, A. and Sjöstrand, J. (2000). Myopisation: the refractive tendency in teenagers. Prevalence of myopia among young teenagers in Sweden. *Acta Ophthalmol Scand*, **78**, 177-181
- Williams, C., Miller, L. L., Gazzard, G. and Saw, S. M. (2008). A comparison of measures of reading and intelligence as risk factors for the development of myopia in a UK cohort of children. *Br J Ophthalmol*, **92**, 1117-1121
- Wong, T. Y., Foster, P. J., Hee, J., Ng, T. P., Tielsch, J. M., Chew, S. J., Johnson, T. J. and Shea, S. K. (2000). Prevalence and risk factors for refractive errors in adult Chinese in Singapore. *Invest Ophthalmol Vis Sci*, **41**, 2486-2494
- Zhao, J., Pan, X., Sui, R., Munos, S. R., Sperduto, R. D. and Ellwein, L. B. (2000). Refractive error in children: results from Shunyi District, China. *Am J Ophthalmol*, **129**, 427-435